

09886197

NEWS 9 Jul 12 BEILSTEIN enhanced with new display and select options,  
resulting in a closer connection to BABS  
NEWS 10 Jul 30 BEILSTEIN on STN workshop to be held August 24 in conjunction  
with the 228th ACS National Meeting  
NEWS 11 AUG 02 IFIPAT/IFIUDB/IFICDB reloaded with new search and display  
fields  
NEWS 12 AUG 02 CAplus and CA patent records enhanced with European and Japan  
Patent Office Classifications  
NEWS 13 AUG 02 STN User Update to be held August 22 in conjunction with the  
228th ACS National Meeting  
NEWS 14 AUG 02 The Analysis Edition of STN Express with Discover!  
(Version 7.01 for Windows) now available  
NEWS 15 AUG 04 Pricing for the Save Answers for SciFinder Wizard within  
STN Express with Discover! will change September 1, 2004

NEWS EXPRESS JULY 30 CURRENT WINDOWS VERSION IS V7.01, CURRENT  
MACINTOSH VERSION IS V6.0c(ENG) AND V6.0Jc(JP),  
AND CURRENT DISCOVER FILE IS DATED 26 APRIL 2004

NEWS HOURS STN Operating Hours Plus Help Desk Availability  
NEWS INTER General Internet Information  
NEWS LOGIN Welcome Banner and News Items  
NEWS PHONE Direct Dial and Telecommunication Network Access to STN  
NEWS WWW CAS World Wide Web Site (general information)

Enter NEWS followed by the item number or name to see news on that  
specific topic.

All use of STN is subject to the provisions of the STN Customer  
agreement. Please note that this agreement limits use to scientific  
research. Use for software development or design or implementation  
of commercial gateways or other similar uses is prohibited and may  
result in loss of user privileges and other penalties.

\* \* \* \* \* STN Columbus \* \* \* \* \*

FILE 'HOME' ENTERED AT 15:23:59 ON 08 AUG 2004

=> FIL STNGUIDE

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	0.21	0.21

FILE 'STNGUIDE' ENTERED AT 15:24:02 ON 08 AUG 2004

USE IS SUBJECT TO THE TERMS OF YOUR CUSTOMER AGREEMENT  
COPYRIGHT (C) 2004 AMERICAN CHEMICAL SOCIETY, JAPAN SCIENCE  
AND TECHNOLOGY CORPORATION, AND FACHINFORMATIONSZENTRUM KARLSRUHE

FILE CONTAINS CURRENT INFORMATION.

LAST RELOADED: Aug 6, 2004 (20040806/UP).

=> file caplus

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	0.06	0.27

FILE 'CAPLUS' ENTERED AT 15:24:26 ON 08 AUG 2004

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2004 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 8 Aug 2004 VOL 141 ISS 7  
FILE LAST UPDATED: 6 Aug 2004 (20040806/ED)

This file contains CAS Registry Numbers for easy and accurate substance identification.

```
=> s isoptera
L1          571 ISOPTERA

=> s l1 and termites
          2133 TERMITES
L2          326 L1 AND TERMITES

=> s coptotermes formosanus
          432 COPTOTERMES
          409 FORMOSANUS
L3          348 COPTOTERMES FORMOSANUS
          (COPTOTERMES(W) FORMOSANUS)

=> s l3 and l2
L4          70 L3 AND L2

=> d ll4 67-70 ibib hitstr abs
'LL4' IS NOT A VALID FORMAT FOR FILE 'CAPLUS'
```

The following are valid formats:

```
ABS ----- GI and AB
ALL ----- BIB, AB, IND, RE
APPS ----- AI, PRAI
BIB ----- AN, plus Bibliographic Data and PI table (default)
CAN ----- List of CA abstract numbers without answer numbers
CBIB ----- AN, plus Compressed Bibliographic Data
DALL ----- ALL, delimited (end of each field identified)
DMAX ----- MAX, delimited for post-processing
FAM ----- AN, PI and PRAI in table, plus Patent Family data
FBIB ----- AN, BIB, plus Patent FAM
IND ----- Indexing data
IPC ----- International Patent Classifications
MAX ----- ALL, plus Patent FAM, RE
PATS ----- PI, SO
SAM ----- CC, SX, TI, ST, IT
SCAN ----- CC, SX, TI, ST, IT (random display, no answer numbers;
              SCAN must be entered on the same line as the DISPLAY,
              e.g., D SCAN or DISPLAY SCAN)
STD ----- BIB, IPC, and NCL
IABS ----- ABS, indented with text labels
```

09886197

IALL ----- ALL, indented with text labels  
IBIB ----- BIB, indented with text labels  
IMAX ----- MAX, indented with text labels  
ISTD ----- STD, indented with text labels

OBIB ----- AN, plus Bibliographic Data (original)  
OIBIB ----- OBIB, indented with text labels

SBIB ----- BIB, no citations  
SIBIB ----- IBIB, no citations

HIT ----- Fields containing hit terms  
HITIND ----- IC, ICA, ICI, NCL, CC and index field (ST and IT)  
                  containing hit terms  
HITRN ----- HIT RN and its text modification  
HITSTR ----- HIT RN, its text modification, its CA index name, and  
                  its structure diagram  
HITSEQ ----- HIT RN, its text modification, its CA index name, its  
                  structure diagram, plus NTE and SEQ fields  
FHITSTR ----- First HIT RN, its text modification, its CA index name, and  
                  its structure diagram  
FHITSEQ ----- First HIT RN, its text modification, its CA index name, its  
                  structure diagram, plus NTE and SEQ fields  
KWIC ----- Hit term plus 20 words on either side  
OCC ----- Number of occurrence of hit term and field in which it occurs

To display a particular field or fields, enter the display field codes. For a list of the display field codes, enter HELP DFIELDS at an arrow prompt (=>). Examples of formats include: TI; TI,AU; BIB,ST; TI,IND; TI,SO. You may specify the format fields in any order and the information will be displayed in the same order as the format specification.

All of the formats (except for SAM, SCAN, HIT, HITIND, HITRN, HITSTR, FHITSTR, HITSEQ, FHITSEQ, KWIC, and OCC) may be used with DISPLAY ACC to view a specified Accession Number.  
ENTER DISPLAY FORMAT (BIB):end

=> d his

(FILE 'HOME' ENTERED AT 15:23:59 ON 08 AUG 2004)

FILE 'STNGUIDE' ENTERED AT 15:24:02 ON 08 AUG 2004

FILE 'CAPLUS' ENTERED AT 15:24:26 ON 08 AUG 2004

L1           571 S ISOPTERA  
L2           326 S L1 AND TERMITES  
L3           348 S COPTOTERMES FORMOSANUS  
L4           70 S L3 AND L2

=> d l4 65-70 ibib hitstr abs

L4   ANSWER 65 OF 70   CAPLUS   COPYRIGHT 2004 ACS on STN  
ACCESSION NUMBER:       1987:529106   CAPLUS  
DOCUMENT NUMBER:       107:129106  
TITLE:                Effect of molybdenum and tungsten compounds on the  
                      survival of **Coptotermes formosanus**  
                      Shiraki (**Isoptera**: Rhinotermitidae) in  
                      laboratory experiments  
AUTHOR(S):            Yoshimura, Tsuyoshi; Tsunoda, Kunio; Nishimoto, Koichi

09886197

CORPORATE SOURCE: Wood Res. Inst., Kyoto Univ., Uji, 611, Japan  
SOURCE: Material und Organismen (1987), 22(1), 47-56  
CODEN: MTOGAF; ISSN: 0025-5270

DOCUMENT TYPE: Journal  
LANGUAGE: English

AB Effects of Mo and W compds. on the termite *C. formosanus* were examined Na molybdate and Na tungstate were effective in diminishing the activity of *C. formosanus*, though the compds. acted very slowly. They caused 100% mortality of *C. formosanus* workers after feeding on 5% treated filter paper for only one day. The slow-action of the compds. may indicate their suitability for the bait-block technique of controlling termite attacks. A remarkable discoloration of the abdomen was observed with **termites** fed on the Na molybdate-treated filter papers and wood blocks.

L4 ANSWER 66 OF 70 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1987:209438 CAPLUS

DOCUMENT NUMBER: 106:209438

TITLE: Characterization of slow-acting insecticides for the remedial control of the Formosan subterranean termite (**Isoptera**: Rhinotermitidae)

AUTHOR(S): Su, Nan Yao; Tamashiro, Minoru; Haverty, Michael I.

CORPORATE SOURCE: Dep. Entomol., Univ. Hawaii, Honolulu, HI, 96822, USA

SOURCE: Journal of Economic Entomology (1987), 80(1), 1-4  
CODEN: JEENAI; ISSN: 0022-0493

DOCUMENT TYPE: Journal

LANGUAGE: English

AB A method is described to exam. time trends in mortality of the Formosan subterranean termite, **Coptotermes formosanus**, exposed to insecticides. Slow-acting toxicants required a longer time to kill **termites** at low concns. than at high concns. The level of mortality and the speed of death were dependent on concentration. With acute toxicants, the time required to kill **termites** was similar at high or low concns., while the mortality levels were concentration-dependent. This speed of death at various concns. of an insecticide can be quantified for comparison purposes using the proposed effective lethal time 90% (ELT90), defined as the amount of time required for an insecticide to kill 90% of the treated individuals within a maximum 14-day span. Slow-acting toxicants were characterized by ELT90 values than spanned a broad range of time, while acute toxicants exhibited a narrow range of ELT90 values.

L4 ANSWER 67 OF 70 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1985:41525 CAPLUS

DOCUMENT NUMBER: 102:41525

TITLE: Evaluation of two insect growth regulators for the bait-block method of subterranean termite (**Isoptera**: Rhinotermitidae) control

AUTHOR(S): Jones, Susan C.

CORPORATE SOURCE: South. Forest Exp. Stn., U.S. Dep. Agric., Gulfport, MS, 39505, USA

SOURCE: Journal of Economic Entomology (1984), 77(5), 1086-91  
CODEN: JEENAI; ISSN: 0022-0493

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The exptl. insect growth regulators fenoxycarb (Ro 13-5223) [72490-01-8] and 2-[p-(m-fluorophenoxy)phenoxy]ethyl ethylcarbamate (Ro 16-1295) [85983-12-6] were effective in the bait-block technique because they caused superfluous intercaste production without adversely affecting feeding of *Reticulitermes virginicus* and **Coptotermes formosanus**. For *R. virginicus*, nos. of nonfunctional intercastes exceeded 50% at 4 wk and survival was significantly reduced at 6 wk. Larvae, workers,

nymphs, and alates of this species developed morphol. abnormalities. At 6 wk, nos. of *C. formosanus* intercastes reached 50%, but significant mortality was not observed. Differences in food substrate altered *C. formosanus* intercaste development; intercastes occurred on treated wood blocks but not on treated  $\alpha$ -cellulose.

L4 ANSWER 68 OF 70 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1984:565481 CAPLUS  
DOCUMENT NUMBER: 101:165481  
TITLE: Structure-activity relationships among aromatic analogs of trail-following pheromone of subterranean **termites**  
AUTHOR(S): Prestwich, Glenn D.; Eng, Waisi; Deaton, Ellen; Wichern, David  
CORPORATE SOURCE: Dep. Chem., State Univ. New York, Stony Brook, NY, 11794, USA  
SOURCE: Journal of Chemical Ecology (1984), 10(8), 1201-17  
CODEN: JCECD8; ISSN: 0098-0331  
DOCUMENT TYPE: Journal  
LANGUAGE: English

AB A series of 12 substituted (Z)-4-phenyl-3-buten-1-ol (PBO) [20047-19-2] derivs. were synthesized and evaluated for trail-following activity in 5 species of subterranean **termites** in the genera *Coptotermes*, *Prorhinotermes*, *Reticulitermes*, and *Schedorhinotermes* (**Isoptera**: *Rhinotermitidae*). The unsubstituted parent PBO was the most active for all species, and electron-withdrawing and electron-donating groups both reduced potency. Sensitivity to substitution in the ortho position suggests steric inhibition of binding by the 2'-substituted analogs. Different sensitivities to these pheromone analogs were found among the 5 species, with *R. flavipes* and *S. lamanianus* showing the highest level of trail-following activity for the PBO analogs.

L4 ANSWER 69 OF 70 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1984:47005 CAPLUS  
DOCUMENT NUMBER: 100:47005  
TITLE: Effects of a dye, Sudan Red 7B, on the Formosan subterranean termite, *Coptotermes formosanus* Shiraki (**Isoptera**: *Rhinotermitidae*)  
AUTHOR(S): Su, Nan Yao; La Fage, Jeffery P.; Esenther, Glenn R.  
CORPORATE SOURCE: Dep. Entomol., Louisiana State Univ., Baton Rouge, LA, 70803, USA  
SOURCE: Material und Organismen (1983), 18(2), 127-33  
CODEN: MTOGAF; ISSN: 0025-5270  
DOCUMENT TYPE: Journal  
LANGUAGE: English

AB Sudan Red 7B [6368-72-5], was selected as a candidate marking material for studying the population dynamics of the Formosan subterranean termite, *C. formosanus*. Its effect on survival and persistence in **termites** was investigated. Almost 100% of the workers that had been allowed to feed on absorbent pads containing 2% (weight/weight) dye for 3 to 9 days and 4% dye for 3 days, retained visible coloration 1 mo after being removed from the source of dye. **Termites** from these treatments also exhibited the lowest mortality, ca. 10%. All **termites** from treatments with higher concns. and/or longer exposure time retained the visible marking 1 mo after the transfer, but, they exhibited higher mortality, i.e. 20-70%. Compared with workers, soldiers exposed to the dye generally exhibited higher mortality than workers and the coloration was less distinctive.

09886197

L4 ANSWER 70 OF 70 CAPLUS COPYRIGHT 2004 ACS on STN  
ACCESSION NUMBER: 1980:210129 CAPLUS  
DOCUMENT NUMBER: 92:210129  
TITLE: Juvenile hormone analogs; effects on the soldier caste  
differentiation in **termites** (  
**Isoptera**)  
AUTHOR(S): Hrdy, Ivan; Krecek, Jan; Zuskova, Zdena  
CORPORATE SOURCE: Prague, Czech.  
SOURCE: Vestnik Ceskoslovenske Spolecnosti Zoologicke (1979),  
43(4), 260-9  
CODEN: VCSZA4; ISSN: 0042-4595  
DOCUMENT TYPE: Journal  
LANGUAGE: English

AB By treating orphaned-grown larvae (or pseudergates) of **termites**  
with juvenile hormones (JHs) or with JH analogs (JHAs), the development of  
presoldier and(or) soldier intercastes was induced. In screening tests  
with *Reticulitermes lucifugus santonensis* and *Prorhinotermes simplex*, JH  
III [22963-93-5] and JHAs hydroprene [41096-46-2], 11-chloro-3,7,11-  
trimethyl-2-dodecenoate [25001-79-0], and tetrahydrofuryl analog of  
methoprene [73618-62-9] were most active. The soldier-caste formation by  
JHs and JHAs was proved in *Kaloterms flavicollis*, *Cryptotermes brevis*,  
*Neotermes castaneus*, *N. jouteli*, *Zootermopsis angusticollis*, *Z.*  
*nevadensis*, *R. lucifugus santonensis*, *P. simplex* and **Coptotermes**  
**formosanus**. The formation of presoldiers was demonstrated in very  
early instars: for *R. lucifugus* from 3rd-instar larvae and for *P. simplex*  
from 2nd-instar larvae. Treatments of starting colonies of *Z. nevadensis*  
and that of colonies of *P. simplex* in natural conditions failed. The use  
of synthetic JHs and JHAs in caste determination and social homeostasis  
studies,  
and the promise of JHAs in social insects control is discussed.

=>  
=> s imidacloprids  
L5 2 IMIDACLOPRIDS

=> s l4 and l5  
L6 0 L4 AND L5

=> s l4 and pests  
8364 PESTS  
L7 1 L4 AND PESTS

=> d 17

L7 ANSWER 1 OF 1 CAPLUS COPYRIGHT 2004 ACS on STN  
AN 2001:423343 CAPLUS  
DN 135:15446  
TI Wood preservatives containing specific plants and insect control of wood  
IN Yoshida, Shinji  
PA Takeda Chemical Industries, Ltd., Japan  
SO Jpn. Kokai Tokkyo Koho, 8 pp.  
CODEN: JKXXAF  
DT Patent  
LA Japanese  
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	JP 2001158009	A2	20010612	JP 1999-342953	19991202

09886197

JP 3326148 B2 20020917  
PRAI JP 1999-342953 19991202

=> s l4 and wood  
145823 WOOD

L8 20 L4 AND WOOD

=> d l8 15-20

L8 ANSWER 15 OF 20 CAPLUS COPYRIGHT 2004 ACS on STN

AN 1993:575821 CAPLUS

DN 119:175821

TI Efficacy of chlorothalonil as a wood preservative against the Formosan subterranean termite

AU Grace, J. Kenneth; Laks, Peter E.; Yamamoto, Robin T.

CS Dep. Entomol., Univ. Hawaii, Honolulu, HI, 96822-2271, USA

SO Forest Products Journal (1993), 43(1), 21-4

CODEN: FPJOAB; ISSN: 0015-7473

DT Journal

LA English

L8 ANSWER 16 OF 20 CAPLUS COPYRIGHT 2004 ACS on STN

AN 1993:75334 CAPLUS

DN 118:75334

TI Termiticidal effects of a glycol borate wood surface treatment

AU Grace, J. Kenneth; Yamamoto, Robin T.

CS Dep. Entomol., Univ. Hawaii, Honolulu, HI, 96822-2271, USA

SO Forest Products Journal (1992), 42(11-12), 46-8

CODEN: FPJOAB; ISSN: 0015-7473

DT Journal

LA English

L8 ANSWER 17 OF 20 CAPLUS COPYRIGHT 2004 ACS on STN

AN 1991:201730 CAPLUS

DN 114:201730

TI Laboratory evaluation of two slow-acting toxicants against Formosan and eastern subterranean **termites** (**Isoptera**: Rhinotermitidae)

AU Su, Nan Yao; Scheffrahn, Rudolf H.

CS Ft. Lauderdale Res. Educ. Cent., Ft. Lauderdale, FL, 33314, USA

SO Journal of Economic Entomology (1991), 84(1), 170-5

CODEN: JEENAI; ISSN: 0022-0493

DT Journal

LA English

L8 ANSWER 18 OF 20 CAPLUS COPYRIGHT 2004 ACS on STN

AN 1988:108090 CAPLUS

DN 108:108090

TI Structure/activity relationships of 2-haloalkanoic acids and their esters as antitermitic agents against Formosan subterranean **termites** (**Isoptera**: Rhinotermitidae)

AU Scheffrahn, Rudolf H.; Su, Nan Yao

CS Inst. Food Agric. Sci., Univ. Florida, Fort Lauderdale, FL, 33314, USA

SO Journal of Economic Entomology (1987), 80(2), 312-16

CODEN: JEENAI; ISSN: 0022-0493

DT Journal

LA English

L8 ANSWER 19 OF 20 CAPLUS COPYRIGHT 2004 ACS on STN

09886197

AN 1987:529106 CAPLUS  
DN 107:129106  
TI Effect of molybdenum and tungsten compounds on the survival of  
**Coptotermes formosanus** Shiraki (**Isoptera**:  
Rhinotermitidae) in laboratory experiments  
AU Yoshimura, Tsuyoshi; Tsunoda, Kunio; Nishimoto, Koichi  
CS Wood Res. Inst., Kyoto Univ., Uji, 611, Japan  
SO Material und Organismen (1987), 22(1), 47-56  
CODEN: MTOGAF; ISSN: 0025-5270  
DT Journal  
LA English

L8 ANSWER 20 OF 20 CAPLUS COPYRIGHT 2004 ACS on STN  
AN 1985:41525 CAPLUS  
DN 102:41525  
TI Evaluation of two insect growth regulators for the bait-block method of  
subterranean termite (**Isoptera**: Rhinotermitidae) control  
AU Jones, Susan C.  
CS South. Forest Exp. Stn., U.S. Dep. Agric., Gulfport, MS, 39505, USA  
SO Journal of Economic Entomology (1984), 77(5), 1086-91  
CODEN: JEENAI; ISSN: 0022-0493  
DT Journal  
LA English

=> d his

(FILE 'HOME' ENTERED AT 15:23:59 ON 08 AUG 2004)

FILE 'STNGUIDE' ENTERED AT 15:24:02 ON 08 AUG 2004

FILE 'CAPLUS' ENTERED AT 15:24:26 ON 08 AUG 2004

L1 571 S ISOPTERA  
L2 326 S L1 AND TERMITES  
L3 348 S COPTOTERMES FORMOSANUS  
L4 70 S L3 AND L2  
L5 2 S IMIDACLOPRIDS  
L6 0 S L4 AND L5  
L7 1 S L4 AND PESTS  
L8 20 S L4 AND WOOD

=> s ll4 and wood

32 LL4  
145823 WOOD

L9 0 LL4 AND WOOD

=> ss l4 and wood

145823 WOOD

L10 20 L4 AND WOOD

=> d l10 14-20 ibib hitsts abs

'HITSTS' IS NOT A VALID FORMAT FOR FILE 'CAPLUS'

The following are valid formats:

ABS ----- GI and AB  
ALL ----- BIB, AB, IND, RE  
APPS ----- AI, PRAI  
BIB ----- AN, plus Bibliographic Data and PI table (default)  
CAN ----- List of CA abstract numbers without answer numbers



09886197

CBIB ----- AN, plus Compressed Bibliographic Data  
DALL ----- ALL, delimited (end of each field identified)  
DMAX ----- MAX, delimited for post-processing  
FAM ----- AN, PI and PRAI in table, plus Patent Family data  
FBIB ----- AN, BIB, plus Patent FAM  
IND ----- Indexing data  
IPC ----- International Patent Classifications  
MAX ----- ALL, plus Patent FAM, RE  
PATS ----- PI, SO  
SAM ----- CC, SX, TI, ST, IT  
SCAN ----- CC, SX, TI, ST, IT (random display, no answer numbers;  
SCAN must be entered on the same line as the DISPLAY,  
e.g., D SCAN or DISPLAY SCAN)  
STD ----- BIB, IPC, and NCL  
  
IABS ----- ABS, indented with text labels  
IALL ----- ALL, indented with text labels  
IBIB ----- BIB, indented with text labels  
IMAX ----- MAX, indented with text labels  
ISTD ----- STD, indented with text labels  
  
OBIB ----- AN, plus Bibliographic Data (original)  
OIBIB ----- OBIB, indented with text labels  
  
SBIB ----- BIB, no citations  
SIBIB ----- IBIB, no citations  
  
HIT ----- Fields containing hit terms  
HITIND ----- IC, ICA, ICI, NCL, CC and index field (ST and IT)  
containing hit terms  
HITRN ----- HIT RN and its text modification  
HITSTR ----- HIT RN, its text modification, its CA index name, and  
its structure diagram  
HITSEQ ----- HIT RN, its text modification, its CA index name, its  
structure diagram, plus NTE and SEQ fields  
FHITSTR ----- First HIT RN, its text modification, its CA index name, and  
its structure diagram  
FHITSEQ ----- First HIT RN, its text modification, its CA index name, its  
structure diagram, plus NTE and SEQ fields  
KWIC ----- Hit term plus 20 words on either side  
OCC ----- Number of occurrence of hit term and field in which it occurs

To display a particular field or fields, enter the display field codes. For a list of the display field codes, enter HELP DFIELDS at an arrow prompt (=>). Examples of formats include: TI; TI,AU; BIB,ST; TI,IND; TI,SO. You may specify the format fields in any order and the information will be displayed in the same order as the format specification.

All of the formats (except for SAM, SCAN, HIT, HITIND, HITRN, HITSTR, FHITSTR, HITSEQ, FHITSEQ, KWIC, and OCC) may be used with DISPLAY ACC to view a specified Accession Number.

ENTER DISPLAY FORMAT (BIB):end

=> d his

(FILE 'HOME' ENTERED AT 15:23:59 ON 08 AUG 2004)

FILE 'STNGUIDE' ENTERED AT 15:24:02 ON 08 AUG 2004

09886197

FILE 'CAPLUS' ENTERED AT 15:24:26 ON 08 AUG 2004

L1 571 S ISOPTERA  
L2 326 S L1 AND TERMITES  
L3 348 S COPTOTERMES FORMOSANUS  
L4 70 S L3 AND L2  
L5 2 S IMIDACLOPRIDS  
L6 0 S L4 AND L5  
L7 1 S L4 AND PESTS  
L8 20 S L4 AND WOOD  
L9 0 S LL4 AND WOOD  
L10 20 SS L4 AND WOOD

=> s l10 15-20 ibib hitstr abs

MISSING OPERATOR L10 15-20

The search profile that was entered contains terms or nested terms that are not separated by a logical operator.

=> d l10 15-20 ibib hitstr abs

L10 ANSWER 15 OF 20 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1993:575821 CAPLUS

DOCUMENT NUMBER: 119:175821

TITLE: Efficacy of chlorothalonil as a wood preservative against the Formosan subterranean termite  
AUTHOR(S): Grace, J. Kenneth; Laks, Peter E.; Yamamoto, Robin T.  
CORPORATE SOURCE: Dep. Entomol., Univ. Hawaii, Honolulu, HI, 96822-2271, USA

SOURCE: Forest Products Journal (1993), 43(1), 21-4

CODEN: FPJOAB; ISSN: 0015-7473

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Chlorothalonil (CTL, tetrachloroisophthalonitrile) was both deterrent and toxic to Formosan subterranean **termites**, **Coptotermes formosanus** (Isoptera: Rhinotermitidae), in laboratory tests using southern yellow pine wafers treated with CTL in oil (AWPA P9 Type A), CTL/chlorpyrifos in oil, or CTL in xylene. The wafers were conditioned by evaporative aging at 40° for 4 wk and exposed to termite attack in a modified ASTM 4-wk (no-choice) test. **Termites** were also exposed to CTL in the xylene carrier and solvent-treated pine wafers in a 4-wk two-choice test for feeding deterrence. CTL retentions were assayed post-test by x-ray fluorescence, and an average 61 % decrease in CTL concentration was found from the pretest nominal retentions. In the no-choice test, CTL retentions of 0.13 to 0.15 pcf (assayed post-test) limited **wood** weight loss from termite feeding to 6-13%, and retentions of 0.26-0.39 pcf CTL resulted in only 3-4% **wood** weight loss. In the two-choice test, CTL retentions ≥0.06 pcf deterred termite feeding in comparison to solvent controls, and the highest tested retention of 0.38 pcf limited **wood** weight loss to 1.5%. Termite mortality was pos. correlated with CTL retention. Assayed CTL retentions ≥0.26 pcf restricted **wood** weight loss from Formosan subterranean termite feeding to <5%. A heavy oil carrier was not essential for CTL efficacy.

L10 ANSWER 16 OF 20 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1993:75334 CAPLUS

DOCUMENT NUMBER: 118:75334

TITLE: Termiticidal effects of a glycol borate **wood** surface treatment

AUTHOR(S): Grace, J. Kenneth; Yamamoto, Robin T.

CORPORATE SOURCE: Dep. Entomol., Univ. Hawaii, Honolulu, HI, 96822-2271,

USA  
 SOURCE: Forest Products Journal (1992), 42(11-12), 46-8  
 CODEN: FPJOAB; ISSN: 0015-7473  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English

AB A remedial wood treatment product known as BORA-CARE, which contains disodium octaborate tetrahydrate (DOT) in a solution of poly- and monoethylene glycols, was evaluated in laboratory tests against the Formosan subterranean termite, *Coptotermes formosanus* (*Isoptera*: Rhinotermitidae). In the first test, hemlock cubes (20 by 20 by 20 mm) were dipped twice in a 1:1 (by volume) aqueous dilution of DOT/glycol (23.54% DOT by weight) and air-dried. All **termites** exposed to the cubes in a laboratory test died within 2 wk, with no feeding on the treated cubes. When a treated cube was placed on top of an untreated cube, **termites** moved over the treated cube, but fed only minimally on the untreated cubes before dying. In the second test, termite feeding and mortality were compared from exposure to **wood** treated with either the DOT/glycol solution or the ethylene glycol solvent for the product. Very limited feeding and 100% termite mortality resulted from exposure to **wood** treated with DOT/glycol. In comparison to the control blocks, treatment with the ethylene glycol solvent alone resulted in a small but significant increase in termite mortality (17%) and decrease in feeding. The high concentration of DOT in poly- and monoethylene glycols deposited on the surface of **wood** treated with DOT/glycol and ingested during termite grooming behavior and/or attempted feeding killed **termites** and protected the **wood** surface from termite penetration.

L10 ANSWER 17 OF 20 CAPLUS COPYRIGHT 2004 ACS on STN  
 ACCESSION NUMBER: 1991:201730 CAPLUS  
 DOCUMENT NUMBER: 114:201730  
 TITLE: Laboratory evaluation of two slow-acting toxicants against Formosan and eastern subterranean **termites** (*Isoptera*: Rhinotermitidae)  
 AUTHOR(S): Su, Nan Yao; Scheffrahn, Rudolf H.  
 CORPORATE SOURCE: Ft. Lauderdale Res. Educ. Cent., Ft. Lauderdale, FL, 33314, USA  
 SOURCE: Journal of Economic Entomology (1991), 84(1), 170-5  
 CODEN: JEENAI; ISSN: 0022-0493  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English

AB Topical toxicity, lethal time, and bait acceptance of two slow-acting toxicants, mirex and sulfluramid, were determined for the Formosan subterranean termite, *Coptotermes formosanus* and the eastern subterranean termite, *Reticulitermes flavipes*. When topically applied to *C. formosanus*, mirex was slightly less toxic (LD50 = 9.14 µg/g) than sulfluramid (LD50 = 6.95 µg/g), but mirex was approx. 34 times more potent (LD50 = 1.78 µg/g) against *R. flavipes* than sulfluramid (LD50 = 60.64 µg/g). Mortality of *R. flavipes* as a function of time was fastest for mirex and slowest for sulfluramid. Lethal time (time to kill 90% of test insects) was similar when *C. formosanus* was treated with either mirex or sulfluramid. Results of a choice bioassay indicated that concentration thresholds of 10 or 30 ppm in **wood** treated with sulfluramid were acceptable to *C. formosanus* and *R. flavipes*, resp. These treatments also produced significant mortality (≥68% mortality at ≥4 ppm for *C. formosanus*, ≥80% mortality at ≥18 ppm for *R. flavipes*) after an 8-wk exposure. **Wood** blocks treated with ≤90 ppm mirex were accepted by *C. formosanus*. Mirex concns.

of  $\geq 10$  ppm produced  $\geq 68\%$  mortality. *R. flavipes* accepted blocks treated with up to 15 ppm of mirex and were killed at significantly higher rates ( $\geq 80\%$ ) when exposed to blocks treated with  $\geq 9$  ppm of mirex.

L10 ANSWER 18 OF 20 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1988:108090 CAPLUS

DOCUMENT NUMBER: 108:108090

TITLE: Structure/activity relationships of 2-haloalkanoic acids and their esters as antitermitic agents against Formosan subterranean **termites** (**Isoptera**: Rhinotermitidae)

AUTHOR(S): Scheffrahn, Rudolf H.; Su, Nan Yao

CORPORATE SOURCE: Inst. Food Agric. Sci., Univ. Florida, Fort Lauderdale, FL, 33314, USA

SOURCE: Journal of Economic Entomology (1987), 80(2), 312-16  
CODEN: JEENAI; ISSN: 0022-0493

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Field-collected Formosan subterranean termite, **Coptotermes formosanus**, foragers were exposed for 2 wk to wood slices containing 5000 ppm of C10-C22 alkanolic and 2-haloalkanoic acids and esters. A feeding reduction index was established to evaluate effects of these compds. on wood consumption by **termites**. **Termites** were maintained for 2 wk after treatment on untreated wood to determine mortality. Unhalogenated acids had minimal effect on *C. formosanus* mortality and wood consumption, but 2-brominated acids were significantly more toxic and diminished feeding. Me esters of haloacids had a variable effect on antitermitic activity that may have been related to carbon-chain length. 2-Iodoctadecanoic acid and ester treatments were more toxic and less fed upon than comparable 2-bromo compds., which, in turn, were more active than their 2-chloro analogs. Methyl-, ethyl-, and isopropyl-2-haloctadecanoates were equally or more toxic than their resp. haloacids.

L10 ANSWER 19 OF 20 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1987:529106 CAPLUS

DOCUMENT NUMBER: 107:129106

TITLE: Effect of molybdenum and tungsten compounds on the survival of **Coptotermes formosanus** Shiraki (**Isoptera**: Rhinotermitidae) in laboratory experiments

AUTHOR(S): Yoshimura, Tsuyoshi; Tsunoda, Kunio; Nishimoto, Koichi

CORPORATE SOURCE: Wood Res. Inst., Kyoto Univ., Uji, 611, Japan

SOURCE: Material und Organismen (1987), 22(1), 47-56  
CODEN: MTOGAF; ISSN: 0025-5270

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Effects of Mo and W compds. on the termite *C. formosanus* were examined. Na molybdate and Na tungstate were effective in diminishing the activity of *C. formosanus*, though the compds. acted very slowly. They caused 100% mortality of *C. formosanus* workers after feeding on 5% treated filter paper for only one day. The slow-action of the compds. may indicate their suitability for the bait-block technique of controlling termite attacks. A remarkable discoloration of the abdomen was observed with **termites** fed on the Na molybdate-treated filter papers and wood blocks.

L10 ANSWER 20 OF 20 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1985:41525 CAPLUS

DOCUMENT NUMBER: 102:41525

09886197

TITLE: Evaluation of two insect growth regulators for the  
bait-block method of subterranean termite (  
**Isoptera**: Rhinotermitidae) control  
AUTHOR(S): Jones, Susan C.  
CORPORATE SOURCE: South. Forest Exp. Stn., U.S. Dep. Agric., Gulfport,  
MS, 39505, USA  
SOURCE: Journal of Economic Entomology (1984), 77(5), 1086-91  
CODEN: JEENAI; ISSN: 0022-0493  
DOCUMENT TYPE: Journal  
LANGUAGE: English  
AB The exptl. insect growth regulators fenoxycarb (Ro 13-5223) [72490-01-8]  
and 2-[p-(m-fluorophenoxy)phenoxy]ethyl ethylcarbamate (Ro 16-1295)  
[85983-12-6] were effective in the bait-block technique because they  
caused superfluous intercaste production without adversely affecting feeding  
of *Reticulitermes virginicus* and *Coptotermes formosanus*  
. For *R. virginicus*, nos. of nonfunctional intercastes exceeded 50% at 4  
wk and survival was significantly reduced at 6 wk. Larvae, workers,  
nymphs, and alates of this species developed morphol. abnormalities. At 6  
wk, nos. of *C. formosanus* intercastes reached 50%, but significant  
mortality was not observed. Differences in food substrate altered *C.*  
*formosanus* intercaste development; intercastes occurred on treated  
wood blocks but not on treated  $\alpha$ -cellulose.

=> s l4 and triazole

21728 TRIAZOLE

L11 0 L4 AND TRIAZOLE

=> logoff

ALL L# QUERIES AND ANSWER SETS ARE DELETED AT LOGOFF

LOGOFF? (Y)/N/HOLD:H

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	67.83	68.10

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE ENTRY	TOTAL SESSION
CA SUBSCRIBER PRICE	-8.82	-8.82

SESSION WILL BE HELD FOR 60 MINUTES  
STN INTERNATIONAL SESSION SUSPENDED AT 15:33:08 ON 08 AUG 2004